

GOLDBACH-LIKE CONJECTURES VIA BERTRAND'S POSTULATE

PETRO KOLOSOV

ABSTRACT. In this manuscript the ternary and binary Goldbach's conjectures are reviewed from Bertrand's postulate prospective. As a result, a relations between both ternary and binary Goldbach's conjectures and Bertrand's postulate are established. Furthermore, a three Goldbach-like conjectures are proposed and discussed. Verification programs are attached to the last section.

CONTENTS

1. Introduction	1
2. Discussion	2
3. Verification	3
References	3

1. INTRODUCTION

In number theory, Bertrand's postulate is a statement that was first conjured in 1845 by Joseph Bertrand [1].

Theorem 1.1. (*Bertrand–Chebyshev theorem.*) *For every positive integer $n > 1$ exists at least one prime p such that*

$$n < p < 2n$$

Bertrand's postulate completely proved by Chebyshev in 1852 [2]. From Bertrand's postulate immediately follows

Lemma 1.1. (*Even Bertrand's partition.*) *By Bertrand's postulate, for every positive integer $n > 1$ there is at least one partition such that*

$$2n = p + \text{odd},$$

where *odd* is odd member of Even Bertrand's partition.

Now we have to recall ternary Goldbach's conjecture or namely Goldbach's weak conjecture

Conjecture 1.2. (*Ternary Goldbach's Conjecture.*) *Every odd number greater than 7 can be expressed as the sum of three odd primes.*

$$2n + 1 = p_i + p_j + p_k, \quad n > 2,$$

where $p_i + p_j + p_k$ is ternary Goldbach partition.

Date: March 13, 2022.

2010 Mathematics Subject Classification. 11P32, 11A41, 97F60.

Key words and phrases. Bertrand's Postulate, Goldbach Conjecture, Ternary Goldbach Conjecture.

Ternary Goldbach's conjecture is claimed to be true by H.A Helfgott [3, 4]. Furthermore, the proof was clarified in [5]. From this prospective, let express the lemma 1.1 for positive odd numbers,

Lemma 1.2. (*Odd Bertrand's partition.*) *By Bertrand's postulate, for every positive integer $n > 1$ there is at least one partition such that*

$$2n + 1 = p + \text{odd} + 1 = p + \text{even},$$

where **even** is even member of Odd Bertrand's partition.

Then we have the following relation between Odd Bertrand's Partition 1.2 and Ternary Goldbach's Partition 1.2, for every positive integer $n > 2$

$$2n + 1 = p + \text{odd} + 1 = \mathfrak{p}_i + \mathfrak{p}_j + \mathfrak{p}_k \quad (1)$$

From equation (1) imply the Goldbach-like conjectures

Conjecture 1.3. *For every prime $p < 2n - 1$ in positive Odd $2n + 1$, $n > 2$ Bertrand's Partition 1.2, the prime p is always a member of Ternary Goldbach's Partition*

$$2n + 1 = \mathfrak{p}_i + \mathfrak{p}_j + p$$

Conjecture 1.4. *For every positive odd $2n + 1$, $n > 2$ Bertrand's partition 1.2, the prime $p < 2n - 1$ always satisfies*

$$\bigvee_{t \in \{i, j, k\}} \mathfrak{p}_t = p,$$

where $\mathfrak{p}_i, \mathfrak{p}_j, \mathfrak{p}_k$ are members of ternary Goldbach's partition of $2n + 1$.

Conjecture 1.5. *For every positive odd $2n + 1$, $n > 2$ Bertrand's partition 1.2 the even part is always a sum of two primes*

$$2n + 1 = p + \text{even} \rightarrow \text{even} = \mathfrak{p}_j + \mathfrak{p}_k, \quad n > 2$$

2. DISCUSSION

Consider the Conjecture 1.3. Suppose it is true. Then for every even integer $2k > 2$, take an odd prime $p < 2k$, and let $n = k + \frac{p-1}{2}$. Then $n < p < 2n - 1$, so the true scenario would imply that there is a ternary Goldbach partition of $2n + 1$ containing p . But $2n + 1 = 2k + p$, giving us a binary Goldbach partition of $2k$.

Suppose the Conjecture 1.3 is false. Then for some $n > 2$, there is a prime p such that $n < p < 2n - 1$ and p is not a member of an odd Goldbach partition of $2n + 1$. It implies that $2n + 1 - p$ is an even integer which is not the sum of two primes.

The Conjecture 1.3 implies Goldbach strong conjecture, however from different prospective. It doesn't assume that all positive even numbers greater than 4 are sum of two primes, but only provides a relation between Bertrand's postulate and ternary Goldbach partition. Conjecture 1.3 is immediately true if Goldbach's strong conjecture is true. Conjectures 1.4, 1.5 are following directly from Conjecture 1.3.

3. VERIFICATION

Conjecture 1.3 may be verified up to 5×10^3 via the program [6]. However, in order to verify larger bounds, the program should be optimized and rewritten using any low-level programming language, for instance, C or C++. Currently, it has an asymptotic complexity of $O(n^2)$ and written on high-level language C#. The verification results are at github.com/kolosovpetro/GoldbachConjecture/PartitionsTo5000.txt

REFERENCES

- [1] J. Bertrand. *Mémoire sur le nombre de valeurs que peut prendre une fonction: quand on y permute les lettres qu'elle renferme*. Bachelier, 1845.
- [2] Tchebichef. Mémoire sur les nombres premiers. *Journal de Mathématiques Pures et Appliquées*, pages 366–390, 1852.
- [3] H. A. Helfgott. Minor arcs for goldbach's problem, 2013. <https://arxiv.org/abs/1305.2897>.
- [4] H. A. Helfgott. The ternary goldbach conjecture is true, 2014. <https://arxiv.org/abs/1312.7748>.
- [5] Harald Andres Helfgott. The ternary goldbach problem, 2015. <https://arxiv.org/abs/1501.05438>.
- [6] Petro Kolosov. Supplementary programs, 2021. <https://github.com/kolosovpetro/GoldbachConjecture>.

Email address: kolosovp94@gmail.com

URL: <https://kolosovpetro.github.io>